

PRESCRIPTION MONOGRAPH

Compounded Active Ingredients: Oxytocin Acetate/PT-141 (Bremelanotide Acetate)

Form: Nasal Spray

Drug Class:

- Oxytocin: Endogenous hypothalamic hormone
 - PT-141: Melanocortin-4 receptor (MC4R) agonist
-

Mechanism of Action:

Oxytocin Acetate is intended to¹:

- Modulate social, emotional, and sexual behavior via central activation of the limbic system and hypothalamus
- Promote sexual receptivity, pair bonding, and arousal
- Enhance parasympathetic tone and smooth muscle relaxation, indirectly supporting erectile function and genital blood flow

PT-141(Bremelanotide) is intended to^{2,3}:

- Activate MC4 receptors in the hypothalamus, which modulate libido and sexual desire
 - Function independently of nitric oxide pathways—making it effective in central sexual dysfunction
 - Increase dopaminergic activity, sexual motivation, and genital arousal in both sexes
-

Indications:

- Hypoactive Sexual Desire Disorder (HSDD) in premenopausal women
 - Low libido or sexual dysfunction in men or women
 - Erectile dysfunction (especially psychogenic)
 - Anorgasmia or delayed orgasm
 - Couples therapy adjunct for intimacy
 - Performance anxiety-related dysfunction
-

Before Use: Let your doctor know if you have had any allergic reactions to nasal sprays in the past. Let your health care provider know if you are pregnant or breast feeding. Let your healthcare provider know of all supplements you are currently taking. Let them know of any thyroid or corticosteroid medications you are prescribed.

Contraindications:

- Pregnancy or lactation (Oxytocin is uterotonic)
 - Uncontrolled hypertension
 - Hypersensitivity to peptides or excipients
 - History of cardiovascular disease
-

Cautions: Let your Healthcare provider know of any changes of vision while on this compounded preparation

How to Use: This medication is a nasal spray and is delivered in a premixed liquid form. Gently blow your nose to clear it from mucous before using the compound. Remove the ring and protective cap to reveal the nasal spray tip. Tilt head slightly forward, hold bottle with thumb on bottom and your index and middle fingers on top. Use your fingers to squeeze down on the pump while you breathe in slowly through your nose. If needed, repeat the process for the second nostril.

PRESCRIPTION MONOGRAPH

Warnings and Precautions:

- PT-141 can increase blood pressure and heart rate; monitor in patients with cardiovascular risk
 - Oxytocin may affect emotional state; psychiatric history should be considered
 - Avoid combining with other serotonergic agents or CNS stimulants unless monitored
-

Adverse Reactions:

Oxytocin:

- Headache
- Emotional lability
- Drowsiness

PT-141:

- Nausea
 - Flushing
 - Transient hypertension
-

Interactions:

Oxytocin:

- SSRIs, SNRIs, MAOIs: Theoretical risk of emotional blunting, mood shifts, or hyponatremia due to overlapping serotonergic effects
- Anxiolytics/sedatives (e.g., benzodiazepines): May enhance sedative effects and emotional sensitivity
- Vasodilators or diuretics: Additive hypotension
- Agents like bupropion or L-DOPA may interact at the emotional/sexual arousal level

PT-141 (Bremelanotide):

- May increase blood pressure and heart rate, counteracting antihypertensive therapies
 - May have reduced efficacy when taken with mu-opioid receptor antagonists like naltrexone
 - Caution is advised with drugs like Tramadol, MDMA, MAOIs due to potential central sympathomimetic synergy and increased risk of hypertension or tachycardia
-

Use in Specific Populations:

- Pregnancy: Contraindicated — Oxytocin has uterotonic effects and may induce uterine contractions. Use is strictly avoided in obstetric settings; PT-141 lacks safety data in pregnancy.
 - Lactation: Caution — Oxytocin is endogenously released during lactation and may stimulate milk letdown; however, safety of exogenous intranasal or combined peptide use during lactation is not established. PT-141 is not recommended.
 - Pediatrics: Not recommended
 - Geriatrics: Permitted with monitoring. Baseline BP and HR should be assessed.
-

Storage:

- Refrigerate at 2°C to 8°C (36°F to 46°F)
 - Can be stored at room temperature (up to 30°C or 86°F) for up to 48hrs
 - Do not freeze
-

Monitoring Parameters:

- Blood pressure (especially with PT-141)
- Sexual response and subjective satisfaction
- Mood and psychological effects
- Frequency of use (tolerance or overuse potential)

PREScription MONOGRAPH

Citations:

1. Ito E, Shima R, Yoshioka T. A novel role of oxytocin: Oxytocin-induced well-being in humans. *Biophys Physicobiol.* 2019;16:132-139. Published 2019 Aug 24. doi:10.2142/biophysico.16.0_132
2. Heiman JR, Kroll R, Liao X, et al. Effects of Bremelanotide (PT-141) on Sexual Function in Women with Hypoactive Sexual Desire Disorder. *J Sex Med.* Published 2021. Available from: <https://kinseyinstitute.org/pdf/JSMedicine-heiman%20et%20al.pdf>.
3. Kingsberg SA, Clayton AH, Portman D, et al. Bremelanotide for the Treatment of Hypoactive Sexual Desire Disorder: Two Randomized Phase 3 Trials. *Obstet Gynecol.* 2019;134(5):899-908. doi:10.1097/AOG.0000000000003500

Compounded medications are not FDA-approved and may differ in risks, benefits, and side effects from FDA-approved products. These statements have not been evaluated by the FDA and are not intended to diagnose, treat or cure any disease or condition and do not indicate any claims of safety or efficacy.
Individual results may vary.